

# The Effect of Chronic Ankle Instability (CAI) on Athletes' Gait and Muscle Activity (EMG): A Systematic Review and Meta-Analysis

Christina Martzivanou<sup>1,\*</sup>, Dimitrios Varvarousis<sup>1</sup>, Dimitrios Dimopoulos<sup>1</sup>, Georgios Ntritsos<sup>2</sup>, Nikolaos Giannakeas<sup>2</sup>, Georgios I. Vasileiadis<sup>1</sup>, Athanasios Kyritsis<sup>1</sup>, Avraam Ploumis<sup>1</sup>

<sup>1</sup>Department of Physical Medicine and Rehabilitation, University of Ioannina, Ioannina, Greece

<sup>2</sup>Department of Informatics and Telecommunications, University of Ioannina, Arta, Greece

**Abstract** Background: The purpose of this study was to evaluate the effect of chronic ankle instability (CAI) on gait and muscle activity (EMG) of athletes. This review focused on the chronic ankle instability (CAI) caused by ankle sprains during athletics. Methods: This review and meta-analysis included 10 studies level I-III: randomized controlled trials (RCT), observational or descriptive laboratory studies and case-control studies. All injured-athletes were compared to healthy controls. Differences in muscle activation between the two groups have been retrieved and documented. Results: Each study demonstrated significant reduction of activation of the affected muscles near the ankle sprain. Overall, the EMG amplitude of peroneus longus (PL), tibialis anterior (TA) and in some cases that of gastrocnemius medius (MG) were decreased after the initial injury, causing the instability of the ankle joint. Conclusion: Despite the differences in how many ankles sprains each athlete has had, all participants revealed significant reduction in muscle activation, specifically that of peroneus longus, thus altering their gait pattern. Significance: The importance of understanding which muscles are activated after an injury is vital not only for post-injured rehabilitation, but furthermore for preventing such injuries and helping young athletes to get back on track on pursuing their athletic careers.

**Keywords** Athletic, Injuries, Electromyography, EMG, Gait, Chronic ankle instability, CAI

## 1. Introduction

Chronic Ankle Instability, shortly known as CAI, is caused by repeated episodes of ankle sprains during sports involving a “dynamic manoeuvre” [1]. According to Doherty et al. [2], a great number of individuals (the number reaching almost half of them) are sustaining an ankle sprain [3] [4]. Besides that, joint instability, pain, swelling, loss of function and gait alteration are chronic residual symptoms often lasting for a life-time [5]. In this systematic review, collected data in the past 20 years have been documented and analysed. The review, consisting of 10 clinical studies reports on information about the participants, their mean age and CAIT score, and their performed tasks. The muscle activation of lower extremity in athletes with chronic ankle instability, after an ankle sprain injury, was measured with the use of attached electrodes. A surface electromyography (sEMG) was placed on the targeted ankle muscles and was compared to the muscular activity of the non-CAI group.

The aim of this study was to evaluate the effect of repetitive ankle sprains on gait and muscle activity (EMG) of athletes. This review focused on the chronic ankle instability (CAI) caused by ankle sprains during athletics. Our purpose was to answer the following questions:

- ✓ What are EMG changes of injured athletes with CAI and how they affect the muscles around the ankle joint?
- ✓ How does the ankle muscular activity of injured athletes with CAI change their gait patterns?

## 2. Methods

### 2.1. Protocol and Registration

In accordance with the PRISMA guidelines [6], our systematic review protocol was registered with the International Prospective Register of Systematic Reviews (PROSPERO) on the 28th of August 2021. The registration number is the following CRD42021270671. The designed study and the noted results were based on the PRISMA statement. Therefore, the review conformed to all PRISMA

\* Corresponding author:

c.martzivanou@gmail.com (Christina Martzivanou)

Received: Oct. 20, 2022; Accepted: Nov. 23, 2022; Published: Dec. 6, 2022

Published online at <http://journal.sapub.org/sports>

guidelines and reported the required information.

## 2.2. Eligibility Criteria

The research was limited to English articles illustrating the effect athletic injuries have on muscle activity (EMG) and gait patterns of athletes. In order to optimize the relevance and accuracy of our research, we decided to include articles published in the last two decades (2001-2021) which were conducted on humans, specifically adults over the age of 19 years. Another important eligible criterion was the type of articles, e.g., randomized controlled trials (RCTs), cross-sectional studies (CSS), case-control studies, meta-analyses and descriptive or observational laboratory studies.

## 2.3. Search Strategy

A computerized literature review was performed including the following databases: MEDLINE, Springer, Research Gate, BMC, Scopus, Cochrane databases. The total number of articles found was 2120 and after filters were applied the number dropped to 418. The last search was conducted on the 15th of December 2021.

## 2.4. Search

The medical search algorithm used in MEDLINE, included following keywords: ("sports" [MeSH Terms] OR "sports" [All Fields] OR "athletic" [All Fields]) OR ("sports" [MeSH Terms] OR "sports" [All Fields] OR "sport" [All Fields]) AND ("injuries" [Subheading] OR "injuries" [All Fields] OR "wounds and injuries" [MeSH Terms] OR ("wounds" [All Fields] AND "injuries" [All Fields]) OR "wounds and injuries" [All Fields]) AND ("electromyography" [MeSH Terms] OR "electromyography" [All Fields] OR "emg" [All Fields]).

## 2.5. Study Selection

The systematic search of the electronic databases (Medline, Springer and PubMed) identified a total of 418 studies, 54 of which were selected for full text screening. The inclusion criteria for participants in these studies were following: adults >19 years, presence of an athletic injury, injury on the lower limbs, participants were athletes (professionals and non-professionals). Studies that mentioned and/or examined underage athletes, non-athletic participants, injury on upper extremities, SCI, tendinopathy, peripheral neuropathy, impairment of the CNS (central nervous system), history of fractures, anatomical lesions (osteochondral), peroneal tendon tears, ligament laxity, ACL ruptures or deficiency were excluded from our study. Only ten studies were considered suitable for data extraction and meta-analysis according to the previous eligibility criteria. The flow chart of the study selection process can be seen in Figure 1.

## 2.6. Data Selection

We extracted individually information on: a) the author and year of publication b) the study-design c) the total number of participants d) the mean age e) the CAIT-score f) which muscles were attached with electrodes g) the data analysis h) undergone tasks by participants and i) the results. Table 1 presents the characteristics of the included studies.

## 2.7. Statistical Analysis

We calculated the summary mean differences, along with the corresponding 95% CI, by pooling the study specific estimates using random-effects models [7]. The presence of heterogeneity was estimated with the Cochran's Q statistic and it was quantified with I [8] [9], when at least 10 studies were included in the meta-analysis. All analyses were performed using Stata (version 14; StataCorp, College Station, TX, [10].

### 2.7.1. Risk Of Bias

The reviewers (CM and AP) documented the methodological quality of the studies and extracted the relevant data. The Biostatistician (GNt) assessed the possible small study effects (an indication of publication bias) by visual inspection of funnel plots and Egger's test, when at least 10 studies were included in the meta-analysis. The PRISMA checklist was applied for the critical appraisal of the studies included, and all articles were examined as full texts.


## 3. Results

### 3.1. Systematic Review

In this systematic review, collected data of 10 clinical studies have been documented and analysed. We focused mainly on prior studies comparing the muscle activity in athletes with chronic ankle instability, after a sequence of ankle sprains, to those with no ankle injury. All injured-athletes were compared to healthy controls. Differences in muscle activation between the two groups have been retrieved and documented. Measurement of lower limb kinetics was performed by using surface electromyography (EMG) on the following muscles: medialis gastrocnemius (MG), rectus femoris (RF), tibialis anterior (TA), peroneus longus (PL), soleus (SOL) and in some cases vastus lateralis (VL), biceps femoris (BF), medial hamstring (MH). The muscle co-contraction index (CCI), which monitors the agonist and antagonist muscles contributing to stabilizing the ankle joint, the Cumberland Ankle Instability Tool (CAIT) and the Identification of Functional Instability Scale (IdFAI) were analysed as well.

**Table 1.** Informational table of the 10 studies, included both in the systematic review as well as in the meta-analysis. Abbreviations: N (Number of participants), TA (tibialis anterior), PL (peroneus longus), MG (medial gastrocnemius), SOL (soleus), RF (rectus femoris), VL (vastus femoris), BiF (biceps femoris), Gmed. (gluteus medius), Gmax. (gluteus maximus), MH (medial hamstring), EMG (Electromyography), CAIT (Cumberland Ankle Instability Tool), IdFAI (Identification of Functional Ankle Instability), FAAM-ADL (Questionnaire of Functional Ankle Ability Measure-Activities of Daily Living), FAAM-sports (Questionnaire of Functional Ankle Ability Measure-Sport Subscale), FAR (Functional Activity Ratio), IC (Initial Contact), CCI (Co-Contraction Index)

Author/Year	Study	N	Age	CAIT	Muscles	Data analysis	Tests	Outcomes
Dejong, 2019	descriptive laboratory	20	21.7 ± 2.32	17.8 ± 4.43	Gmax and Gmed	USI, CAIT, IdFAI, FAAM-ADL, FAAM-Sport	gait cycle (stance and swing phase)	Gmed: decrease of CAIT, FAR measures, FAAM-ADL and FAAM-Sport scores. Increased IdFAI scores and kinesiophobia. No significant differences in Gmax.
Delahunt, 2007	case-control	26/50 had FI	25.6 ± 6.1	n/m	TA, RF, PL, SOL	3D kinematics and EMG during lateral hop task for the period 200 ms pre- and post-IC	force plate-standing, 10 lateral hops	Significant increase of iEMG amplitude (pre- IC) in TA, SOL
Koldenhoven, 2016	case-control laboratory	17/34 with CAI	20 ± 2.6	n/m	TA, PL, MG, Gmed	sEMG (RMS areas and amplitudes), plantar pressure (peak pressure, pressure-time integral, time to peak pressure, contact area and time, location of COP)	gait cycle (stance and swing phase)	CAI group: laterally deviated COP (stance phase) and corresponding increases in peak pressure. Significantly lower sEMG amplitude in TA and higher sEMG in PL, MG and Gmed.
Li, 2018	Pair-matched control	21/42 with CAI	21 ± 2	19.3 ± 6	TA, RF, PL, GL, VL, BiF	SEMG amplitude, GRF, CCR, CCI	Drop landing	Pre-landing: less ankle stability (lower CAIT and greater IdFAI scores), lower PL and greater GL activation, significantly greater ankle muscle co-contraction in the frontal plane. Landing phase: greater activations for TA, RF, VL
Lin, 2011	Controlled laboratory	15/30 with CAI	21.6 ± 2.4	18.1 ± 5	TA, PL, GL	EMG amplitude, GRF, CI of TA/PL (frontal plane) and TA/GL (sagittal plane), ankle kinematics (dorsiflexion-plantar flexion angle, inversion-eversion angle and internal-external rotation angle)	Running and stop-jump landing	Running (pre-landing): significantly higher ankle inversion angle and lower dynamic ankle joint stiffness. Stop-jump landing (post-landing): greater ankle inversion and lower peak ankle eversion angle
Nanbancha, 2019	Matched group	19	20.58 ± 1.54	18.47 ± 3.67	MG, PL, TA	EMG amplitude, ankle dorsiflexion, plantar flexion, inversion and eversion, MEP (Motor Evoked Potential), latency	Jump-landing	Lower CAIT score and MEP concerning TA and PL. Longer latency duration concerning TA and MG. Lower activity of PL.
Santilli, 2005	Descriptive laboratory	14	26.4 ± 3.99	n/m	PL	SEMG amplitude	Gait cycle (stance and swing phase)	Significant decrease in PL muscle activity.
Simpson, 2019	Controlled laboratory	15/30 with CAI	21.3 ± 1.6	18.9 ± 3.7	TA, MG, PL and PB	EMG, CCI and ankle kinematics (expected and unexpected ankle inversion perturbations; inversion angle at IC, time to max. inversion angle, max. inversion angle and velocity)	Single leg drop-landing	Greater maximum inversion angle, less inversion angle at IC, less TA muscle activity and frontal plane CCI 200 ms post-landing, and prolonged latency of the PB.
Son, 2017	Controlled laboratory	22/66 with CAI	22.7 ± 2	n/m	TA, PL, MG, Gmax, Gmed, VL, MH	EMG amplitude	Jump-landing/cutting	Decrease in plantar flexion, inversion and hip abduction angle. Reduction in TA, PL, MG, VL, Gmed and Gmax muscle activity. Increase in knee and hip flexion angle.
Tretriluxana, 2020	Laboratory cross-sectional	20/40 with CAI	20.6 ± 1.5	18.6 ± 3.7	TA, PL, MG	Statistic parametric mapping (SPM) 200 msec. before foot-contact with the ground during a jump-landing task and EMG amplitude	Jump- landing	Decreased EMG of PL. The critical threshold was 12.1 and the supra-threshold cluster was 65% prior to foot contact with the ground for the CAI group.



**Figure 1.** Flowchart [11] of the process of gathering the published relative literature

Table 1 presents the collected information of each study, ordered by the year of publication. Data such as type of study, number of participants, mean age, measured EMG amplitude of specific muscles, Cumberland Ankle Instability Tool (CAIT) score and the task/test that participants had to complete in order to assess the muscular activation, have each been collected and summarised. The participants were in their early to mid-twenties, stated a CAIT- score of <19 and most of them were asked to perform the stop, jump and landing test. The muscle activation of CAI and non-CAI athletes was measured either by surface electromyography (sEMG) and ankle kinematics (dorsiflexion–plantar flexion angle, inversion-eversion angle and internal-external rotation angle) or motor evoked potential (MEP), evident in the study of Nanbancha *et al* [12].

### 3.1.1. Results of Systematic Review

The first article, that of DeJong *et al* [13], showed a decrease of CAIT of gluteus medius, decreased FAR measures, FAAM-ADL and FAAM-Sport scores. The idFAI scores were increased and no significant differences were seen in the gluteus maximus muscle. The CAI-athletes revealed a kinesiophobia after all. In the study of Delahunt *et al* [14], a significant increase of iEMG amplitude (pre- IC) in tibialis anterior and soleus was noticed.

The study of Koldenhoven [15] on the other hand, resulted in laterally deviated COP (stance phase) and corresponding increases in peak pressure. Significantly lower sEMG amplitude of tibialis anterior and higher sEMG of peroneus longus, gastrocnemius medialis, gluteus medius.

Li *et al* [16] presented following results in the pre-landing phase: less ankle stability (lower CAIT and greater IdFAI scores), lower muscle activation of peroneus longus, greater gastrocnemius lateralis activation and significantly greater ankle muscle co-contraction in the frontal plane. In the

landing phase the muscle activation of tibialis anterior, rectus femoris and vastus longus was greater.

Moreover, Lin *et al* [17] showed that in the running task (pre-landing phase) the ankle inversion angle was significantly higher with a lower dynamic ankle joint stiffness. The stop-jump landing test (post landing) revealed a greater ankle inversion and lower peak ankle eversion angle.

In addition, Nanbancha *et al* presented lower CAIT score and MEP concerning tibialis anterior and peroneus longus, longer latency duration concerning tibialis anterior and gastrocnemius medialis. A significant lower muscle activity of peroneus longus in athletes with CAI was displayed as well, something that was common in the study of Santilli *et al* [18].

Simpson and his co-authors [19] demonstrated a greater maximum inversion angle in athletes with chronic ankle instability, less inversion angle at initial contact (IC), less muscle activity of the tibialis anterior and lastly a prolonged latency of the peroneus brevis muscle.

Additionally, Son *et al* [20], found a decrease in plantar flexion, inversion and hip abduction angle, a reduction in muscle activity of the tibialis anterior, peroneus longus, vastus longus, gluteus maximus and medius, gastrocnemius medialis and finally an increase in knee and hip flexion angle. The last study being that of Tretriluxana *et al* [21], displayed a decreased electromyography of the muscle peroneus longus.

In general, the pre-initial contact (200 ms prior) tasks appeared to have:

- Decreased electromyography of peroneus longus, tibialis anterior, gastrocnemius medialis, vastus longus and gluteus maximus and medius
- Decrease in plantar flexion, inversion and hip abduction angle
- Increased knee and hip flexion angle
- Prolonged latency of tibialis anterior and gastrocnemius medialis

As opposed to the post initial contact (landing phase) tests where following was evident:

- Slower dynamic ankle joint stiffness
- Prolonged latency of peroneus brevis
- Greater ankle inversion
- Lower peak ankle eversion angle

### 3.2. Meta-Analysis


A meta-analysis was executed on the following 4 outcomes as presented on Table 2 (see Table 2): Cumberland Ankle Instability Tool score (CAIT), Identification of Functional Ankle Instability score (idFAI), co-contraction index (CCI) in sagittal and frontal plane and lastly the Questionnaire of Functional Ankle Ability Measure-Sport Subscale score (FAAM-sport).

**Table 2.** Data collection of CAIT score, idFAI, CCI in sagittal and frontal plane, FAAM-Sports.

	N	Gender	Age	SD	CAIT	SD	idFAI	SD	FAAM - sport	SD	CCI (sag.)	SD	CCI (fro.)	SD
Dejong, 2019														
CAI	20	10 M, 10 F	21.7	2.32	17.8	4.4 3	21.2	4.05	72.83	5.5 6				
Control	20	10 M, 10 F	21.2	2.8	30	0	1	1	100	0				
Delahun, 2007														
CAI	26	16 M, 10 F	25.6	6.1							0.58	0.26	0.77	0.36
Control	24	15 M, 9 F	22.6	4.3							0.61	0.38	0.52	0.35
Koldenhoven, 2016														
CAI	17	6 M, 11 F	20	2.6			21.3	5.2	75	6.9				
Control	17	6 M, 11 F	21.8	4.3			N/a	N/a	100	0				
Li, 2018														
CAI	21	21 F	21	2	19.3	6	22.2	9.2			52.5	20.4	57.8	15
Control	21	21 F	21	2	29.5	0.9	1.3	2.1			51.7	15.9	67.5	18
Lin, 2011														
CAI	15	6 M, 9 F	21.6	2.4	18.1	5								
Control	15	7 M, 8 F	21.5	2.6	30	0								
Nanbanacha, 2019														
CAI	19	15 M, 4 F	20.58	1.54	18.47	1.74								
Control	19	15 M, 4 F	20.58	1.3	29.11	1.05								
Santilli, 2005														
CAI	14	10 M, 4 F	26.4	3.99										
Simpson, 2019														
CAI	15		21.3	1.6	18.9	3.7					0.53	0.25	0.66	0.22
Control	15		21.5	1.5	29.7	0.6					0.48	0.23	0.66	0.23
Son, 2017														
CAI	20	12 M, 8 F	22.7	2					60.9	11.6				
Control	20	12 M, 8 F	21.8	2.3					100					
Tretriluxana, 2020														
CAI	20	16 M, 4 F	20.6	1.5	18.6	3.7					76.29	17.45	54.77	14.78
Control	20	16 M, 4 F	20.5	1.3	29.2	1					62.07	16.68	45.85	13.61


The CAIT score was significantly lower in athletes with chronic ankle instability in compare to non-CAI athletes. The mean difference (MD) of 4 out of 10 studies in total led to this significance ( $I^2= 88.7\%$ ,  $p = 0.000$ ). In the study of Li

et al the mean difference (95% CI) was  $-2.38 (-3.18, -1.58)$ . The mean difference of Nanbanacha was  $-7.40 (-9.23, -5.58)$ . Simpson and Tretriluxana scored with  $-4.07 (-5.36, -2.79)$  and  $-3.91 (-4.99, -2.84)$  respectively (Figure 2).




**Figure 2.** Data synthesis for the studies reporting on CAIT score. Athletes with CAI showed significant lower CAIT scores compared to the non-CAI group. The mean difference with a 95% CI and the weight each study had are also depicted

While the CCI in both sagittal ( $I^2= 38.7\%$ ,  $p = 0.180$ ) and frontal plane ( $I^2= 67.4\%$ ,  $p = 0.027$ ) showed no statistical significance (Figure 3 and 4), the idFAI score appeared to be increased in athletes with CAI ( $I^2= 93.2\%$ ,  $p = 0.000$ ).




**Figure 3.** Data synthesis of the 4 studies measuring the CCI in sagittal plane, however with no statistical significance seen



**Figure 4.** Data synthesis of the 4 studies measuring the CCI in frontal plane; with no statistical significance as well

The statistical importance of the increased idFAI score is rather small in contrast to the other outcomes, as the meta-analysis of idFAI itself was conducted on two studies only, that of Li and DeJong (Figure 5).



**Figure 5.** Data synthesis of the idFAI score. Although the idFAI was significantly higher in CAI-athletes, the statistical importance of this meta-analysis is poor, considering the small sample size (2 studies)

## 4. Discussion

The purpose of this study was to evaluate the effect of CAI on gait analysis and muscle activity of athletes. As anticipated, athletes with chronic ankle instability (CAI), displayed different muscular activation in regard to non-CAI athletes. In our meta-analysis, athletes with CAI showed significant lower CAIT scores compared to the non-CAI group. The mean difference with a 95% CI was -4.30 (-6.04; -2.56). The idFAI was significantly higher in CAI-athletes with a mean difference and 95% CI of 4.92 (1.28; 8.56).

Due to alterations in biomechanics of lower extremities during high-impact movements (i.e., movements often seen in athletic manoeuvres), the chronic ankle instability shows different neuromuscular control when compared to athletes with no injuries.

In the study of Lin *et al.*, this difference is described as a "deficit in feedback neuromuscular control", where the central nervous system adjustment procedure is delayed. The reason behind the deficient feedback in neuromuscular control being the inability of the ankle to execute the motion normally hence remaining inverted. In addition, the muscles around the ankle joint seem to adapt for maintaining the joint stability in athletes with CAI, through decreasing the activation of the tibialis anterior muscle (TA).

Thus said, less ankle stability (lower CAIT and greater IdFAI scores) and lower maximum voluntary isometric contraction (MVIC) moments for ankle dorsiflexors and evertors are documented in athletes with chronic ankle instability. In the pre-landing phase, 200 milliseconds prior to initial foot contact, CAI-athletes displayed altered muscle activation when compared to non-CAI athletes. The EMG demonstrated reduced peroneal longus activation and greater

activation of gluteus lateralis. In similar studies, peroneal, gluteus maximus and medius strength-deficits are frequently reported for CAI-patients. A tendency of increased vastus lateralis-activation has also been noticed. The ankle muscle co-contraction was therefore significantly greater in the frontal plane.

Differences in muscle activation of the ankle joint were also present in the landing phase. The EMG captured a greater activation of the following muscles: tibialis anterior, rectus femoris and vastus lateralis. There was a tendency of reduced biceps femoris activation and the co-contraction index (CCI) of the ankle muscles was greater in sagittal as well as frontal planes. In our meta-analysis however, we had no statistical significance in both frontal and sagittal planes. Moreover, knee muscle activations such as increase in knee extensor motions and co-contraction ratio (CCR) CAI patients adopted landing positions of less inversion, less plantarflexion, more knee and hip flexion, and less hip abduction angle during the initial contact to mid-landing phase (0%–25% of stance) compared with copers and controls.

While increased activation of the tibialis anterior in athletes with CAI could increase the ankle stability in the landing, it could also hinder the ability of ankle energy absorption and further influence the knee biomechanics and muscle activations. Therefore, the changes in ankle muscle stimulation may result in some atypical knee muscle loadings, such as greater co-contraction ratio (CCR) of quadriceps to hamstrings, which was confirmed in our study too.

## 5. Study Limitations

There were some limitations to the present study. On the one hand, we have an all-female study which may limit the generalizability of the results to female athletes with CAI and on the other hand they are two studies without any gender-information. Our meta-analysis was limited by the small sample size of the studies. The results could possibly lack of power to adequately justify our findings. There is a need for more research on this topic, with studies that will have a larger sample of patients and use modern technologies to measure gait parameters and muscle activity. Therefore, our results should be considered with a mild degree of alertness in regards to the sample size.

## 6. Conclusions

In conclusion, the results of this study confirm altered muscle activation in the lower extremities and different gait pattern in athletes with Chronic Ankle Instability (CAI) when compared to the control groups (coppers). In general, a significant lower CAIT score by CAI athletes was noticed in each of the ten studies relative to non-CAI athletes. Alterations such as reduced peroneal longus activation in the prelanding phase and increased co-contraction of ankle

muscles in the landing phase were noticed in athletes with CAI [22]. The modified muscle movements, whether they are reduced or increased, in athletics during exercises may act as a protecting and stabilizing mechanism. Some suggestions involved exercises that stabilize and strengthen the lower limb muscles or proximal musculature during gait-training, but there is a need for more research on this topic, with studies that will have a larger sample of patients and use modern technologies to measure gait parameters and muscle activity.

## Conflicts of Interest

None

## Funding

The project MEGATRON (Megadata analysis of robot-assisted base and 3D camera system for improved kinetic rehabilitation - subproject 1: Implementation and Evaluation - MIS 5047227) is co-funded by the European Union and the Hellenic Republic.

---

## REFERENCES

- [1] Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train.* 2007; 42:311.
- [2] Doherty C, Delahunt E, Caulfield B, Hertel J, Ryan J, Bleakley C. The incidence and prevalence of ankle sprain injury: a systematic review and meta-analysis of prospective epidemiological studies. *Sports Med.* 2014; 44:123-140.
- [3] Gribble PA, Delahunt E, Bleakley C, et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the International Ankle Consortium. *Br J Sports Med.* 2014; 48:1014-1018.
- [4] Anandacoomarasamy A, Barnsley L. Long term outcomes of inversion ankle injuries. *Br J Sports Med.* 2005; 39(3): 14.
- [5] Gribble, P.a., Robinson, R.H., 2009. An examination of ankle, knee, and hip torque production in individuals with chronic ankle instability. *J. Strength Cond. Res.* 23, 395–400. <http://dx.doi.org/10.1519/JSC.0b013e3181818efbb2>.
- [6] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71. Doi: 10.1136/bmj.n71.
- [7] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials.* 1986;7(3): 177-88.
- [8] Kulinskaya E, Dollinger MB. An accurate test for homogeneity of odds ratios based on Cochran's Q-statistic. *BMC Med Res Methodol.* 2015; 15:49.
- [9] Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ.* 1997

- Sep 13; 315(7109): 629-34. Doi: 10.1136/bmj.315.7109.629. PMID: 9310563; PMCID: PMC2127453.
- [10] StataCorp. 2015. Stata Statistical Software: Release 14. College Station, TX: StataCorp LP.
- [11] Moher et al, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7): e1000097. doi:10.1371/journal.pmed1000097.
- [12] Nanbancha A, Tretriluxana J, Limroongreungrat W, Sinsurin K. Decreased supraspinal control and neuromuscular function controlling the ankle joint in athletes with chronic ankle instability. *Eur J Appl Physiol*. 2019 Sep; 119(9): 2041-2052. Doi: 10.1007/s00421-019-04191-w. Epub 2019 Jul 18. PMID: 31321512.
- [13] DeJong AF, Mangum LC, Hertel J. Gluteus medius activity during gait is altered in individuals with chronic ankle instability: An ultrasound imaging study. *Gait Posture*. 2019 Jun; 71:7-13. Doi: 10.1016/j.gaitpost.2019.04.007. Epub 2019 Apr 11. PMID: 30999270.
- [14] Delahunt E, Monaghan K, Caulfield B. Ankle function during hopping in subjects with functional instability of the ankle joint. *Scand J Med Sci Sports*. 2007 Dec; 17(6): 641-8. Doi: 10.1111/j.1600-0838.2006.00612.x. Epub 2007 Mar 1. PMID: 17331087.
- [15] Koldenhoven, R.M., Feger, M.A., Fraser, J.J. et al. Surface electromyography and plantar pressure during walking in young adults with chronic ankle instability. *Knee Surg Sports Traumatol Arthrosc* 24, 1060–1070 (2016). <https://doi.org/10.1007/s00167-016-4015-3>.
- [16] Yumeng Li, Jupil Ko, Marika A. Walker, Cathleen N. Brown, Julianne D. Schmidt, Seock-Ho Kim, Kathy J. Simpson. Does chronic ankle instability influence lower extremity muscle activation of females during landing. *Journal of Electromyography and Kinesiology*. Volume 38, 2018. Pages 81-87. ISSN1050-6411. <https://doi.org/10.1016/j.jelekin.2017.11.009>.
- [17] Lin C-F, Chen C-Y, Lin C-W. Dynamic Ankle Control in Athletes with Ankle Instability During Sports Maneuvers. *The American Journal of Sports Medicine*. 2011; 39(9): 2007-2015. Doi:10.1177/0363546511406868.
- [18] Santilli V, Frascarelli MA, Paoloni M, et al. Peroneus Longus Muscle Activation Pattern during Gait Cycle in Athletes Affected by Functional Ankle Instability: A Surface Electromyographic Study. *The American Journal of Sports Medicine*. 2005; 33(8): 1183-1187.
- [19] Jeffrey D. Simpson, Ethan M. Stewart, Alana J. Turner, David M. Macias, Samuel J. Wilson, Harish Chander, Adam C. Knight. Neuromuscular control in individuals with chronic ankle instability: A comparison of unexpected and expected ankle inversion perturbations during a single leg drop-landing. *Human Movement Science*, Volume 64, 2019. Pages 133-141, ISSN 0167-9457.
- [20] SON, S. J., H. KIM, M. K. SEELEY, and J. T. HOPKINS. Movement Strategies among Groups of Chronic Ankle Instability, Coper, and Control. *Med. Sci. Sports Exerc.*, Vol. 49, No. 8, pp. 1649–1661, 2017.
- [21] Tretriluxana J, Nanbancha A, Sinsurin K, Limroongreungrat W, Wang HK. Neuromuscular control of the ankle during pre-landing in athletes with chronic ankle instability: Insights from statistical parametric mapping and muscle co-contraction analysis. *Phys Ther Sport*. 2021 Jan; 47:46-52. Doi: 10.1016/j.ptsp.2020.11.023. Epub 2020 Nov 4. Erratum in: *Phys Ther Sport*. 2020 Nov 22; PMID: 33166739.
- [22] Gribble PA, Bleakley CM, Caulfield BM, et al. Evidence reviews for the 2016 international ankle consortium consensus statement on the prevalence, impact and long-term consequences of lateral ankle sprains. *Br J Sports Med*. 2016; 50: 1496-1505.